



*Selective broadleaf weed herbicide
Specially formulated for control of broadleaf weeds in turf[^] (parks, lawns,
golf courses, cemeteries, sports fields, and other grassy areas.)*

**Kills over 60 listed broadleaf weeds
without harming lawns[^]**

Low-Odor Formulation

Microtechnology[†] improves penetration of reduced active ingredient formulation.

[^]see back label for listed weeds and control details.

[^]Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

[†]Microtechnology signifies emulsion particles less than 1 micron.

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt*.....	0.885%
Mecoprop-p, dimethylamine salt**.....	0.573%
Dicamba, dimethylamine salt***.....	0.109%

OTHER INGREDIENTS:^{††}

Total 100.000%

*2,4-D Acid Equivalent	0.738%
** Mecoprop-p Acid Equivalent	0.478%
*** Dicamba Acid Equivalent	0.091%

^{††}Contains Petroleum Distillate

KEEP OUT OF REACH OF CHILDREN

Shake well before using

Protect from Freezing

See inside booklet for additional Precautionary Statements and Directions for Use

MADE IN CANADA (MAY CONTAIN IMPORTED PARTS) FOR:

Intelligro™

2310 Lakeshore Road West
Mississauga, Ontario, Canada L5J 1K2
1-844-388-2665

Patent(s): www.suncor.com/patents

intelligro™

EPA EST. NO. 79693-CAN-001
EPA REG. NO. 69526-15
641021/3

[™]Trademark of Suncor Energy Inc. Used under license.

ESL121815 REV022519

PRODUCT SUMMARY

CONTROLS: Kills over 60 broadleaf weeds without harming lawns.^

^Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

TREATMENT AREA : 1 fl.oz. product treats up to 172 ft²

MICROTECHNOLOGY :

Fast-acting, improved absorption formula.

Patent-pending formulation improves herbicide absorption into weeds. Improved penetration reduces amount of active ingredients required in the formula to kill weeds.



HOW IT WORKS

CIVITAS WEEDfree BRAND Concentrate kills over 60 broadleaf weeds, without harming your lawn.^

Patent-pending formula improves the penetration of active ingredients into the weed.

^Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

FIRST AID

If swallowed:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. To reach a U.S. POISON CONTROL CENTER, call 1-800-222-1222. You may also contact INTELLIGRO: 1-844-388-2665, or CANUTEC Canadian Transport Emergency Centre: 613-996-6666.

NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard, contains petroleum distillate.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are Barrier Laminate, Nitrile rubber, Butyl, Neoprene rubber or Viton. If you want more options, follow the instructions for Category E on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves (except for applicators using groundboom equipment), when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate, and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking or ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Read and follow all label directions.

Protect from freezing

Shake well before using

AGRICULTURAL USE REQUIREMENTS

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeve shirt and short pants,
- chemical-resistant footwear plus socks,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant headgear for overhead exposure,
- protective eyewear.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

USE PRECAUTIONS



Do not apply as a fine mist because of potential injury to desirable plants. Do not use this product on or near desirable plants, or on the exposed roots of desirable trees and shrubs, injury may result.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. People or pets can enter treated area after sprays have dried.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, airblast, and chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles. When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, beans, cotton, flowers, fruit trees (foliage), grapes (in growing stage), okra, ornamentals, soybeans (vegetative stage), sunflowers, tobacco, tomatoes, and other vegetables. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Do not spray exposed roots of ornamentals and trees.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for ground boom applications:

Do not apply with a nozzle height greater than 4 feet above the canopy.

Precaution on warm season Grasses: Centipede, St. Augustine, Bahia, Common Bermuda, and Zoysia

Some temporary discoloration may occur on warm season grasses. Do not apply if grass is stressed from heat or drought. Avoid application during transition periods from/to dormancy. **Centipede and St. Augustine** grasses are sensitive to 2,4-D and great care should be taken to prevent overdosing. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent discoloration or injury. Also, applications to dormant Bermudagrass, dormant Zoysia grass, and dormant Bahiagrass are not suggested.

- **IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

ADDITIONAL RESTRICTIONS

- Do not mix with liquid fertilizers that contain nitrates.
- Do not apply through any type of irrigation system.
- Do not apply near lakes, streams, rivers ponds, or ditchbanks.
- Do not apply by air.
- Do not use more product than specified on the label.
- Do not apply when the lawn, turf or site is under severe stress from drought, insect attack or heat.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants.
- Do not pour spray solutions near these plants.
- Do not apply in windy conditions.
- Do not irrigate within 24 hours of application.

MAXIMUM APPLICATIONS PER YEAR	APPLICATION MAXIMUM	APPLICATION INTERVAL
Limited to 2 applications per treatment area per year.	The maximum rate (8 gal./acre) for a single application is 0.75 lb ae/acre (0.9 lb DMA salt/acre) (0.424 lb 2,4-D ae, 0.274 lb MCPP-p ae, 0.052 lb Dicamba ae per acre per application).	Repeat application if needed, but with no less than 30 days between applications.

APPLICATION DIRECTIONS

WEEDS CONTROLLED IN TURF or LAWN AND ORNAMENTAL AREAS:

Bedstraw	Grass leaved stitchwort	Russian thistle
Bindweed (field)	Ground ivy (creeping Charlie)	Sandwort (thyme-leaved) [†]
Birdsfoot trefoil [†]	Hawkweed	Sheep sorrel
Black medic	Heal-all	Shepherd's purse
Blackseed plantain	Henbit	Smartweed (green)
Blueweed	Honeysuckle [†]	Soliva
Brambles [†]	Jimsonweed	Sow thistle
Bull thistle	Knotweed	Speedwell (purslane leaved)
Burdock	Kochia	Spotted spurge
Buttercup	Kudzu	Spreadwell (annual)
Canada thistle	Lambsquarters	Spurge [†]
Carolina geranium	Little starwort	Stitchwort [†]
Chickweed	Mallow	Stonecrop (mossy)
Chicory	Morningglory	Sumac [†]
Clover (White clover in California)	Mouse-ear chickweed	Veronica (thyme-leaved) [†]
Cocklebur	Mustards	Vervain [†]
Common chickweed	Narrow-leaved plantain	Vetch [†]
Cudweed	Oxalis (yellow woodsorrel)	Violet
Daisy fleabane	Ox-eye daisy	Wild aster
Dandelion	Pennywort	Wild blackberry [†]
Devil's paint brush (hawkweed)	Peppergrass	Wild carrot
Dock (curled)	Pigweed	Wild garlic
English daisy	Pineapple weed [†]	Wild geranium
Evening primrose [†]	Plantains (narrow-leaved, buckhorn, broadleaf)	Wild lettuce
Fall dandelion (fall hawkbit)	Poison ivy	Wild onion
False dandelion	Poorjoe	Wild radish
Fleabane	Povertyweed [†]	Wild raspberry [†]
Florida pusley	Prostrate pigweed	Yarrow
Frenchweed poison oak	Purslane	Yellow hawkweed
	Ragweed	

[†]Not registered for use in California

**TURF AND ORNAMENTAL (GOLF COURSES, CEMETERIES, PARKS,
SPORT FIELDS, TURFGRASS, LAWNS AND OTHER GRASS AREAS)
(Post emergence)**

WHERE TO USE

This product is for use as a broadcast or spot treatment on lawns for broadleaf weed control without damaging lawn grasses.

IMPORTANT: Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass.

NOTE: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

WHEN TO USE

- Do not water within 24 hours after spraying.
- Do not apply when rain or irrigation is anticipated.
- Spray when weeds are young and actively growing. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing.
- Apply when daytime temperatures are above 45°F and below 85°F.
- Newly seeded lawns may be treated after the third mowing.
- If the lawn and/or turf is dry, water 24 hours before using this product.
- Grass seed can be sown 3 to 4 weeks after application.
- Delay mowing 1 to 2 days before and after the application of this product.

HOW TO USE

For broadcast treatment:

For smaller units: Apply 0.25 - 2.75 gals. of product per 1000 sq ft with the specified amount of water. For larger units: Apply 2 - 8 gallons of product per acre with the specified amount of water. Table below provides the recommended product rate and spray volume. Mix well before use. Use the lower rate on annual or small weeds. For best control of large weeds and tough perennial weeds, such as dandelion, use one application of the higher product rate or two applications of the lower rate making the second treatment 30 days following the first application

Broadcast treatment

Step 1 - You must know and calibrate your sprayer carrier volume.

Step 2 - Use this chart to determine product rate based on the known calibrated carrier volume.

Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Smaller Units – 1000 SQ FT	Step 1 - Carrier Volume	Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Larger Units - Acre
Rate / 1000 sqft.	Carrier Volume / 1000 sq. ft.	Carrier Volume / Acre
6 fl. oz.	0.25-0.35 gal.	10 to 15 gal.
9 fl. oz.	0.35-0.45 gal.	15 to 20 gal.
12 fl. oz.	0.45-1.15 gal.	20 to 50 gal.
15 to 18 fl. oz.	1.15-1.85 gal.	50 to 80 gal.
18 to 24 fl. oz.	1.85-2.75 gal.	80 to 120 gal.
		Rate / Acre
		2 gal.
		3 gal.
		4 gal.
		5-6 gal.
		6-8 gal.

Avoid overlapping between spray patterns.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

California

Broadcast treatment

Step 1 - You must know and calibrate your sprayer carrier volume.

Step 2 - Use this chart to determine product rate based on the known calibrated carrier volume.

Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Smaller Units – 1000 SQ FT	Step 1 - Carrier Volume		Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Larger Units - Acre
Rate / 1000 sqft.	Carrier Volume / 1000 sq. ft.	Carrier Volume / Acre	Rate / Acre
12 fl. oz.	0.45-1.15 gal.	20 to 50 gal.	4 gal.
15 to 18 fl. oz.	1.15-1.85 gal.	50 to 80 gal.	5-6 gal.
18 to 24 fl. oz.	1.85-2.75 gal.	80 to 120 gal.	6-8 gal.

For spot treatment using hand-operated sprayers:

Mix 10 fl.oz. of the product with 118 fl.oz. (3.7 qts.) of water to make 1 gallon of total spray solution. Hold sprayer about 12 inches from weed and spray to lightly wet weeds.

TURF, GROWN FOR SEED OR SOD (Post emergence)

WHERE TO USE

For selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

IMPORTANT: Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

WHEN TO USE

- Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 24 hours after application.
- Spray when weeds are young and actively growing. Apply when daytime temperatures are above 45°F and below 85°F.
- If the sod is dry, water 24 hours before using this product.
- Delay mowing 1 to 2 days before and after the application of this product.
- Treated areas may be reseeded 3 to 4 weeks after application.

HOW TO USE

For broadcast treatment:

For smaller units: Apply 0.25 - 2.75 gals. of product per 1000 sq ft with the specified amount of water. For larger units: Apply 2 - 8 gallons of product per acre with the specified amount of water. Table below provides the recommended product rate and spray volume. Mix well before use. Use the lower rate on annual or small weeds. For best control of large weeds and tough perennial weeds, such as dandelion, use one application of the higher product rate or two applications of the lower rate making the second treatment 30 days following the first application

Broadcast treatment

Step 1 - You must know and calibrate your sprayer carrier volume.

Step 2 - Use this chart to determine product rate based on the known calibrated carrier volume.

Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Smaller Units – 1000 SQ FT	Step 1 - Carrier Volume		Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Larger Units - Acre
Rate / 1000 sqft.	Carrier Volume / 1000 sq. ft.	Carrier Volume / Acre	Rate / Acre
6 fl. oz.	0.25-0.35 gal.	10 to 15 gal.	2 gal.
9 fl. oz.	0.35-0.45 gal.	15 to 20 gal.	3 gal.
12 fl. oz.	0.45-1.15 gal.	20 to 50 gal.	4 gal.
15 to 18 fl. oz.	1.15-1.85 gal.	50 to 80 gal.	5-6 gal.
18 to 24 fl. oz.	1.85-2.75 gal.	80 to 120 gal.	6-8 gal.

Avoid overlapping between spray patterns.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages

California

For California, apply only 4-8 gallons/acre, as follows:

Broadcast treatment

Step 1 - You must know and calibrate your sprayer carrier volume.

Step 2 - Use this chart to determine product rate based on the known calibrated carrier volume.

Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Smaller Units - 1000 SQ FT	◀	Step 1 - Carrier Volume		▶	Step 2 - CIVITAS WEEDfree BRAND Concentrate Rate Larger Units - Acre
Rate / 1000 sqft.	◀	Carrier Volume / 1000 sq. ft.	Carrier Volume / Acre	▶	Rate / Acre
12 fl. oz.	◀	0.45-1.15 gal.	20 to 50 gal.	▶	4 gal.
15 to 18 fl. oz.	◀	1.15-1.85 gal.	50 to 80 gal.	▶	5-6 gal.
18 to 24 fl. oz.	◀	1.85-2.75 gal.	80 to 120 gal.	▶	6-8 gal.

For spot treatment using hand-operated sprayers:

Mix 10 fl.oz. of the product with 118 fl.oz. (3.7 qts.) of water to make 1 gallon of total spray solution. Hold sprayer about 12 inches from weed and spray to lightly wet weeds.

NEWLY SEEDED AREAS: Applying this product to grass seedlings is recommended after the third mowing.

NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS: Delay applications to newly sodded, sprigged, or plugged grasses 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysia grass, and dormant Bahiagrass are not suggested.

EXPECTED RESULTS



Visible injury on weeds can be seen within 12 hours. It can take up to 4 weeks for weeds to completely die.

How often to apply:

Repeat application can be made after 30 days if necessary. The maximum number of broadcast applications per treatment site is 2 per year.

STORAGE AND DISPOSAL



Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE:

Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place in a locked storage area.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons.

Confine spills. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL:

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

Rigid, nonrefillable containers, equal to or less than five gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Rigid nonrefillable containers, greater than 5 gallons:

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non-rigid, nonrefillable container (275-gallon bladder with frame):

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Return frame to dealer.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Intelligro, the manufacturer or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law neither Intelligro, the manufacturer, nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product. To the extent consistent with applicable law or except as warranted by this label, Intelligro, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and Intelligro, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.